




Each tablet contains 525 mg FEBANTEL, 504 mg PYRANTEL EMBONATE and 175 mg PRAZIQUANTEL.

Tablets for the treatment and control of

- Roundworm (Ascarids)
- Hookworm
- Whipworm
- Tapeworm (including *Taenia ovis*)

And for the control of the shedding of cysts of the protozoan *Giardia* spp. in dogs.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

TREATMENT PROGRAMMES FOR WORMS

Roundworms: *Toxocara canis*, *Toxascaris leonina*

Hookworms: *Ancylostoma caninum*, *Uncinaria stenocephala*

Whipworm: *Trichuris vulpis*

Tapeworms: *Taenia* spp., *Dipylidium caninum*

Puppies: Puppies should be treated every 2 weeks from the age of two to twelve weeks, then monthly until 6 months of age.

Adult Dogs: (above 6 months of age) should be dewormed every 3 months. Dogs heavily infested with roundworms, hookworms or whipworms may require retreatment after 14 days. Where tapeworm control only is required, for example where flea tapeworm segments keep reappearing 2-3 weeks after treatment due to reinfestation, or where *Taenia ovis* control is required, use DRONCIT between DRONTAL TASTY BONE 35 kg treatments.

TREATMENT PROGRAMMES FOR GIARDIA

Giardia infection: Treat for three consecutive days at recommended dose. Do not exceed recommended doses.

USE DURING PREGNANCY AND LACTATION

The safety of DRONTAL TASTY BONE 35 kg has not been investigated during the 1st and 2nd part of pregnancy, and is therefore not recommended during these periods. When treating for worms, it is recommended to provide single treatment during the 3rd part of pregnancy, 10 days prior to whelping.

The product may be used during lactation.

Use of DRONTAL TASTY BONE 35 kg for the 3 days treatment against *Giardia* spp. infections in the 3rd part of pregnancy should be based on a benefit/risk assessment conducted by a veterinarian.

DOSAGE AND ADMINISTRATION

Special precautions to be taken by the person administering the veterinary medicinal product to animals.

In the interests of good hygiene, persons administering the tablet directly to the dog or by adding it to the dog’s food, should wash their hands afterwards.

In case of *Giardia* infection treatment of the dog: clean thoroughly and disinfect or move the dogs whilst on treatment, particularly in kennel situations. *Giardia* could infect human beings so please ask for medical advice in case your dog is infected.

DRONTAL TASTY BONE 35 kg is given as a single dose by mouth. The tablets are flavoured and may be administered with or without food. Access to normal diet does not need to be limited before or after treatment.

The dose rate is 1 tablet per 35 kg (15 mg febantel, 14.4 mg pyrantel embonate and 5 mg praziquantel/kg body weight).

DOSE RATE FOR ORAL ADMINISTRATION	
Body weight (kg)	Tablet quantity
7 – 17.5	½
>17.5 – 35	1
>35 – 52.5	1 ½
>52.5 – 70	2

For each additional 17.5 kg body weight, administer an additional half tablet. Not for use in dogs weighing less than 7 kg.

Interaction with other medicinal products

Do not use with other veterinary medicines containing the ingredient piperazine.

Overdose

Up to 10 times the recommended dose of the product was tolerated without problems in dogs and puppies.

Contraindications

Do not use in case of hypersensitivity to the active substances or to any of the excipients.

Adverse Reactions

In very rare cases, mild short term digestive tract effects (eg. vomiting) may occur.

Special Warning

Fleas serve as intermediate hosts for one common type of tapeworm- *Dipylidium caninum*. Tapeworm infestation is certain to re-occur unless control of intermediate hosts such as fleas, mice etc. is undertaken.

READ ENTIRE LABEL BEFORE USE

May cause an allergic skin reaction. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May damage fertility or the unborn child.



HANDLING PRECAUTIONS

Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Wear protective gloves, protective clothing, eye protection, face protection. Wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace. Store locked up.

FIRST AID

IF exposed or concerned: Get medical advice or attention. IF INHALED: Remove person to fresh air and keep comfortable for breathing. If experiencing respiratory symptoms: Call a POISON CENTER or doctor. IF ON SKIN: Wash with plenty of water. If skin irritation or rash occurs: Get medical advice or attention. Take off contaminated clothing and wash it before reuse. For advice contact the National Poisons Centre 0800 POISON (0800 764 766).

ENVIRONMENTAL PROTECTION

Toxic to aquatic life with long lasting effects. Avoid release to the environment. Collect spillage.



EMERGENCY PROTECTION

Place in an appropriate waste disposal container. For specialist advice in an emergency call CHEMTREC 0800 425 459. If additional hazard information is required, refer to Safety Data Sheet. For a copy please go to www.elanco.co.nz.

DISPOSAL

Preferably dispose of the product by use. Dispose of empty packaging by wrapping in paper and placing in garbage.

STORAGE

Store below 30 °C. After opening the blister, remaining half tablets should be returned to the open blister. Once opened, use left-over half-tablets by 7 days.

Drontal Tasty Bone 35 kg is registered pursuant to the ACVM Act 1997, No. A011600. See www.foodsafety.govt.nz for registration conditions. Droncit ACVM No. A002919

Elanco New Zealand
106 Wiri Station Road, Manukau, Auckland 2104
Customer Info Line: 0800 446 121
www.mypetandi.elanco.com/nz

Drontal, Droncit, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.
© 2025 Elanco or its affiliates